

LISTING OF THE CLAIMS

1. (Currently amended) A method ~~to purify~~ for purifying crude diacerein ~~by the toluene approach, wherein the diacerein is obtained by acetylation and chrome oxidation,~~ characterized by comprising the following steps:

- a) ~~crude diacerein is dissolved~~ dissolving the crude diacerein in acetone/water[[,]];;
- b) ~~adjusting the pH is adjusted~~ adjusting the pH with a tertiary amine in acetone[[,]];;
- c) ~~stirring the solution is stirred for about six hours~~ stirring the solution[[,]];;
- d) ~~adding an organic water immiscible-solvent is added and~~ adding an organic water immiscible-solvent ~~is added and~~ to the solution is stirred,;
- e) ~~separating the organic solvent phase is separated~~ separating the organic solvent phase from the acetone/water phase[[,]];;
- f) ~~the extraction is repeated~~ repeating steps d) and e) about 5 to 15 times ~~with the organic water immiscible solvent and, every time, the organic phase is separated from the acetone/water phase~~,;
- g) ~~crystallizing the diacerein is crystallized~~ crystallizing the diacerein from the acetone/water phase ~~upon by~~ by changing the pH from neutral to acid with a strong acid[[,]]; and
- h) ~~collecting the crystallized product is centrifuged or filtered, washed with water and dried~~ collecting the crystallized product ~~is centrifuged or filtered, washed with water and dried~~ by centrifugation or filtration.

2. (Currently amended) A method ~~to purify~~ for purifying crude diacerein ~~by the toluene approach~~ The method of claim 1, wherein ~~the diacerein is obtained by acetylation and chrome oxidation,~~ according to ~~claim 1~~, characterized in that in step a), the crude diacerein is dissolved, while wet, in a 1/1 acetone/water mixture in an proportion of about 13 times the volume of dry diacerein, and it is stirred until completely dissolved.

3. (Currently amended) A method ~~to purify~~ for purifying crude diacerein ~~by the toluene approach~~ The method of claim 1, wherein ~~the diacerein is obtained by acetylation and chrome oxidation,~~ according to ~~claim 2~~, characterized in that in step b), the pH is adjusted at to 6.6-7.2, preferably 7.0-7.2, with a solution of a tertiary amine in acetone.

4. (Currently amended) ~~A method to purify crude diacerein by the toluene approach, wherein the diacerein is obtained by acetylation and chrome oxidation, according to The method of claim 3, characterized in that a wherein the tertiary amine is a trialkylamine with C1-4 alkyl groups may be used as a tertiary amine, preferably selected from the group: trimethylamine, triethylamine, tripropylamine, methyldiethylamine, diethylpropylamine, among which triethylamine is most preferred.~~

5. (Currently amended) ~~A method to purify crude diacerein by the toluene approach, wherein the diacerein is obtained by acetylation and chrome oxidation, according to The method of claim 2 1, characterized in that wherein in step c), stirring lasts the solution is stirred for about six hours in order for the diacerein to dissolve completely.~~

6 (Currently amended) ~~A method to purify crude diacerein by the toluene approach, wherein the diacerein is obtained by acetylation and chrome oxidation, according to The method of claim 1, characterized in that wherein in step d), once the diacerein is completely dissolved, it is purified by organic extraction with a the water-immiscible[[-]]solvent , which is selected from the group consisting of[[:]] benzene, toluene, xylene, isobutyl acetate and ethyl acetate, among which toluene is most preferred.~~

7. (Currently amended) ~~A method to purify crude diacerein by the toluene approach, wherein the diacerein is obtained by acetylation and chrome oxidation, according to The method of claim 1, characterized in that wherein in step e) the extraction of diacerein with the organic water immiscible-solvent is performed[[:]] in a volume of toluene organic solvent equivalent to 1.6 volumes of the dry diacerein is added to the dissolution; the dissolution is stirred and the organic phase is separated from the acetone/water solution; about 5 to 15 repeated extractions (step f) are performed with the same amount of organic water immiscible solvent and with the same proportion of the solvent to dry diacerein.~~

8. (Currently amended) ~~A method to purify crude diacerein by the toluene approach, wherein the diacerein is obtained by acetylation and chrome oxidation, according to claim 7,~~

~~characterized in that a minimum of 10 repeated extractions is preferred~~ The method of claim 1,
wherein steps d) and e) are performed between 10 and 15 times.

9. (Currently amended) ~~A method to purify crude diacerein by the toluene approach,~~
~~wherein the diacerein is obtained by acetylation and chrome oxidation, according to claim 7,~~
~~characterized in that about 15 repeated extractions with the organic solvent are preferably~~
~~performed~~ The method of claim 8, wherein steps d) and e) are performed about 15 times.

10. (Currently amended) ~~A method to purify crude diacerein by the toluene approach,~~
~~wherein the diacerein is obtained by acetylation and chrome oxidation, according to~~ The method
of claim 1, ~~characterized in that~~ wherein in step g), after at least 10 extractions with the organic
~~water immiscible solvent,~~ the diacerein is crystallized in water at a pH between about 2.5 and
about 3.0 with a strong acid, such as sulphuric, hydrochloric or phosphoric acid.

11. (Currently amended) ~~A method to purify crude diacerein by the toluene approach,~~
~~wherein the diacerein is obtained by acetylation and chrome oxidation, according to~~ The method
of claim 10, ~~characterized in that 80% phosphoric acid is preferred~~ wherein the strong acid is
selected from the group consisting of sulfuric acid, hydrochloric acid and phosphoric acid..

12. (Currently amended) ~~A method to purify crude diacerein by the toluene approach,~~
~~wherein the diacerein is obtained by acetylation and chrome oxidation, according to~~ The method
of claim 1, ~~characterized in that in step h), it is~~ further comprising washing the centrifuged or
filtered crystallized product with water and ~~dried~~ drying said washed product.

13. (Currently amended) ~~A method to purify crude diacerein by the toluene approach,~~
~~wherein the diacerein is obtained by acetylation and chrome oxidation, according to~~ The method
of claim 1, ~~characterized in that~~ wherein the yield of diacerein is obtained under a 90-93% yield,
~~with an~~ the average purity of is 99.17%, ~~a~~ the content of aloe-emodin ~~of is~~ is about 7-10 ppm and ~~a~~
the content of chromium of about 20-25 ppm.

14. (New) The method of claim 3, wherein the pH is adjusted to 7.0-7.2.

15. (New) The method of claim 4, wherein the tertiary amine is selected from the group consisting of trimethylamine, triethylamine, tripropylamine, methyldiethylamine and diethylpropylamine.

16. (New) The method of claim 6, wherein the water-immiscible solvent is toluene.

17. (New) The method of claim 11, wherein the strong acid is phosphoric acid.